

## ARTÍCULO/ARTICLE

**Nesting biology of an Ecuadorian endemic hummingbird, the endangered Violet-throated Metaltail *Metallura baroni***Agustín Carrasco-Ugalde<sup>1</sup>, Paul Molina Abril<sup>1</sup>, Daniel Pacheco<sup>1</sup>, Boris A. Tinoco<sup>2,\*</sup><sup>1</sup> Autilloproducciones<sup>2</sup> Escuela de Biología, Universidad del Azuay, Av. 24 de Mayo 7-77, Cuenca, Ecuador.\*Corresponding author, e-mail: [btinoco@uazuay.edu.ec](mailto:btinoco@uazuay.edu.ec)

Editado por/Edited by: Juan Freile

Recibido/Received: 28 Mayo 2021 Aceptado/Accepted: 14 Junio 2022

Publicado en línea/Published online: 18 Julio 2022

**Biología de anidación de un colibrí endémico del Ecuador, el amenazado Metalura Gorjivioleta *Metallura baroni*****Resumen**

El colibrí Metalura Gorjivioleta *Metallura baroni* es una especie amenazada, endémica de los Andes del sur del Ecuador. La biología reproductiva de esta especie era casi desconocida. En este artículo describimos el nido, huevos y desarrollo de pichones de esta especie en base al seguimiento de nueve nidos encontrados en la vía Cuenca-Molleturo-Naranjal, en el Parque Nacional Cajas, durante 2019. Detectamos un periodo reproductivo entre marzo y mayo, que corresponden a la estación lluviosa. Los nidos fueron encontrados en taludes al borde de la vía, y fueron construidos con musgos, ramas pequeñas, raicillas, corteza de árboles de *Polylepis* sp. (Rosaceae) y fibras de *Puya* sp. (Bromeliaceae). En todos los nidos encontramos dos huevos ovales blancos. Los huevos y pichones fueron atendidos exclusivamente por la hembra. El tiempo desde la eclosión de los huevos hasta abandono del nido fue de 28–32 días. Anidar en el borde de la vía podría ser riesgoso debido a la probabilidad de atropellamiento por vehículos; por tanto, estudiar el éxito reproductivo de *M. baroni* en los nidos localizados en el borde de la vía es una prioridad de investigación para la especie. La información presentada en este artículo incrementa el conocimiento sobre la historia natural de *M. baroni*.

**Palabras clave:** Parque Nacional Cajas, trampa ecológica, colibrí, páramo, reproducción.**Abstract**

Violet-throated Metaltail *Metallura baroni* is a threatened hummingbird species, endemic to the Andes of southern Ecuador. Details about the reproductive biology of this species are largely unknown. In this manuscript, we describe the nest, eggs, and nestlings' development based on nine nests found along with the Cuenca-Molleturo-Naranjal road, in Cajas National Park, in 2019. We document a breeding period from March to May, which corresponds to the rainy season. All nests were found in roadside embankments, and were constructed with mosses, small twigs, rootlets, *Polylepis* sp. (Rosaceae) bark, and *Puya* sp. (Bromeliaceae) fibers. In all nests, we found two white oval eggs that were exclusively attended by a female across the breeding period. Hatching to fledging period lasted 28–32 days. Nesting at roadsides can be particularly risky due to a high probability of collision with vehicles; thus, an assessment of the breeding success of *M. baroni* on nests located at roadsides is a research priority for this species. The information provided here increases the knowledge about the natural history of *M. baroni*.

**Keywords:** Cajas National Park, ecological trap, hummingbird, páramo, reproduction.**INTRODUCTION**

Violet-throated Metaltail *Metallura baroni* is a hummingbird that belongs to the *Metallura aenocauda* clade (Graves, 1980; García-Moreno, 1999; Benham *et al.*, 2015) in the monophyletic genus *Metallura*. It is endemic to the provinces of Cañar and Azuay, in the western Andes of southern Ecuador. *Metallura baroni* ranges from

3000 to 4000 m a.s.l. (Tinoco *et al.*, 2009; Astudillo *et al.*, 2015), occupying mainly shrubby páramo and *Polylepis* woodland, where it feeds on several short corolla flowers, including *Brachyotum* spp. (Melastomataceae), *Berberis* spp. (Berberidaceae), and *Barnadesia arborea* (Asteraceae) (Tinoco *et al.*, 2009). As a result of its small distribution range and fragmented populations that are declining in numbers over time, *M. baroni* is considered as Endangered, both at national (Freile *et al.*, 2019) and global levels (BirdLife International, 2021).

Most of the ecological knowledge about *M. baroni* comes from studies focused on its distribution, general behavior and habitat use (Tinoco *et al.*, 2009; Astudillo *et al.*, 2014, 2015). However, its breeding biology remains largely unknown (Schuchmann, 1999; Tinoco *et al.*, 2009). The only breeding information available to date comes in a brief report of a single, cup-shaped nest that contained a single egg, found in April in Cajas National Park, southern Ecuador (Schuchmann, 1999).

Many traits related to the breeding biology of bird species tend to show high levels of phylogenetic conservatism (Zyskowski & Prum, 1999; Fang *et al.*, 2018); therefore, generalities about the breeding biology of *Metallura* species might inform about the reproduction of *M. baroni*. *Metallura* nests are usually located not high above the ground, mainly in rock ledges, caves, and embankments (Snow 1980; Schuchmann, 1999; Mamani-Cabana, 2020; Matta-Pereira *et al.*, 2019; Soto-Patiño *et al.*, 2021). Common nesting materials include mosses, lichens, small bird feathers, and mammal fur. As with other hummingbirds, females lay two eggs, and nest attendance is provided exclusively by the female (Schuchmann, 1999). However, it remains to be determined if those generalities apply to *M. baroni*.

Knowledge about the breeding biology of a species is key for learning about its natural history and it also can be useful for guiding scientifically sound conservation actions. In this manuscript, we provide detailed information about several aspects of the breeding biology of *M. baroni*, including characteristics of the nests, eggs, incubation time, and development of nestlings. This manuscript increases the knowledge about the basic biology of this endangered species, and the information presented can guide further research on the factors that influence the breeding success of the species.

## METHODS

### Study area

This study was conducted in Cajas National Park (CNP), Azuay province, southern Andes of Ecuador (-2.787222, -79.203055). This is a high-Andean national park with an elevation range from 3100–4450 m a.s.l. (Rodríguez-Girón *et al.*, 2014). The temperature in the area has extreme daily fluctuations, commonly varying from 18 C° during day hours to -2 C° at night; the total annual precipitation ranges between 1000–1800 mm (Céleri *et al.*, 2007), presenting a bimodal pattern with increasing rains from March to May, and from October to December (Campozano *et al.*, 2016). The main vegetation type in CNP is páramo grasslands, with montane shrubs and montane forest occurring at lower elevations (below 3500 m a.s.l.). CNP is considered an important bird area for conservation (Freile & Santander, 2005; Astudillo *et al.*, 2015), and it is part of the core area of Macizo del Cajas Biosphere Reserve (Rodríguez-Girón *et al.*, 2014).

### Data collection

We searched for nests of *M. baroni* along the borders of the Cuenca-Molleturo-Naranjal road, between Control Quinuas, 25.8 km from Cuenca (-2.784649, -79.193071; 3650 m a.s.l.) and Tres Cruces, 35.6 km from Cuenca (-2.777661, -79.240735; 4150 m a.s.l.). This is a paved road located in the north part of CNP, and connects two main cities in Ecuador, Cuenca and Guayaquil.

From January through February 2019, we visited the study area once every two weeks searching for nests, but when breeding activity was detected, we increased our sampling effort to visits every 3–6 days, until late May, when we did not find any new nests. On every visit, two observers walked along the two sides of the road, placing particular attention on embankments and rocky walls, as those are common places used by *Metallura* species to nest (Schuchmann, 1999; Mamani-Cabana, 2020; Soto-Patiño *et al.*, 2021). The walks always started from Tres Cruces, at 8h00–10h00, and lasted c. 2–3 h. We are aware that roadsides might not be a high-quality habitat for *M. baroni* (Astudillo *et al.*, 2014; Tinoco *et al.*, 2019), but given the descriptive aims of this study, our observations are useful to provide basic ecological knowledge for the species.

The identity of nests was always confirmed by the observation of an adult *M. baroni* attending the nest. For each nest, we recorded the following variables: geographical coordinates with a handheld GPS, main habitat type around the nest, and nest shape type (*sensu* Simon & Pacheco, 2005). If the nest was located at less than 2 m in height above the ground, we took measurements of its external and internal diameter, height and depth of the internal chamber; all these measurements were taken with an analog caliper. We also noted the height above the ground of the nest using a measuring tape. For nests located above 2 m in height, we only estimated height above the ground in meters, and checked the status of the nest by using photos taken with a camera attached to an aluminum pole. One of the nests was collected after the breeding period to describe the materials used in its construction. Every nest found was visited at least once every 6 days to document the outcome of the breeding effort.

To obtain data about incubation time and nest attendance, we installed a time-lapse PlotWatcher Pro camera (Day 6 Outdoors) in one nest, once during the incubation period, for a total of 509 min, and once during the nestling period, at day 20 after hatching, for 233 min. The camera was set to take one picture every second. We could not install cameras in the other nests because most of them were exposed and located on the border of the road. We described the development of a pair of nestlings in one nest that was visually monitored by the observers from the hatchling day until nest abandonment by fledglings. We created a feather development table of the nestlings following Jongsomjit *et al.*, (2007).

## RESULTS

### Nest locations and season

We found a total of nine nests, all located between 3520–3990 m a.s.l. (Table 1). The first active nest was found on 4 March, while the last new active nest was found on 14 April. Eight nests were found in overhanging banks located at the border of the road (Fig. 1A), and one nest was found on the border of a small creek (less than 1 m wide), 5 m away from the road. In all cases, the surrounding habitat was páramo grasslands.

Table 1: General information about nine nests of Violet-throated Metaltail *Metallura baroni* found along the Cuenca-Molleturo-Naranjal road, Azuay province, Ecuador, in 2019.

| Nest ID | Geographic Coordinates   | Elevation (m a.s.l.) | Encounter date | Nesting stage in the encounter date | History of the nest                         |
|---------|--------------------------|----------------------|----------------|-------------------------------------|---|
| 1       | -2.800348,<br>-79.29603  | 3525                 | 4 March 2019   | 2 nestlings                         | 1 egg depredated; 1 fledgling left the nest |
| 2       | -2.790281,<br>-79.202500 | 3885                 | 11 March 2019  | 2 eggs                              | 2 eggs depredated                           |
| 3       | -2.787222,<br>-79.203055 | 3820                 | 16 March 2019  | 2 nestlings                         | 2 fledglings left the nest                  |
| 4       | -2.789444,<br>-79.201388 | 3875                 | 22 March 2019  | 2 nestlings                         | Unknown                                     |
| 5       | -2.787222,<br>-79.206666 | 3925                 | 22 March 2019  | 2 nestlings                         | 2 nestlings died in the nest                |
| 6       | -2.783888,<br>-79.223333 | 3980                 | 22 March 2019  | 2 eggs                              | 2 eggs hatched, no further monitoring       |
| 7       | -2.782777,<br>-79.188611 | 3630                 | 27 March 2019  | 2 nestlings                         | 1 egg depredated; 1 fledgling left the nest |
| 8       | -2.788242,<br>-79.205513 | 3925                 | 4 April 2019   | 2 eggs                              | 2 eggs depredated                           |
| 9       | -2.784166,<br>-79.205555 | 3990                 | 14 April 2019  | 2 nestlings                         | 2 fledglings left the nest                  |



Figure 1: A) Location of a Violet-throated Metaltail *Metallura baroni* nest (red circle) along an embankment in the Cuenca-Molleturo-Naranjal road, in a section that crosses Cajas National Park (Boris Tinoco). B) Placement of a *M. baroni* nest (red arrow) in the top of an embankment of the Cuenca-Molleturo-Naranjal road (Boris Tinoco).

### Nest descriptions

The nests were found at 3.61 m above the ground, on average ( $N = 8$ ;  $SD = 1.87$  cm; range = 0.4–6.03 cm). The nests were attached to small rock ledges or were constructed within overhanging thin roots (Fig. 1B). The shape of all nests was open cup/late type. We took detail measurements of four nests, which presented the following mean sizes: external diameter 7.86 cm ( $SD = 0.87$  cm; range = 7.36–9.25 cm), height 8.64 cm (range = 5.23–12.77 cm), internal diameter 4.13 cm ( $SD = 0.78$  cm; range = 3.42–4.97 cm), depth of the internal chamber 3.28 cm ( $SD = 0.59$  cm; range = 2.71–4.1 cm).

The outer layer of the one nest we disaggregated was made out of mosses, dry twigs, small rootlets, *Polylepis* (Rosaceae) bark, small dry leaves, spiderweb, animal fur, and small white flowers (Fig. 2A). The inner chamber was made of soft *Puya* sp. (Bromeliaceae) downy fibers and small feathers (Fig. 2B). Similar materials were observed in the other nests.



Figure 2: A) Shape and materials of a Violet-throated Metaltail *Metallura baroni* nest; an adult female incubating is present (Paúl Molina). B) Eggs (Boris Tinoco).

### Eggs and nestlings

All the nests contained two white, oval eggs (Fig. 2B). Nestlings hatched with pink skin, yellow bill, and light cream neossotiles in their dorsum (Fig. 3A). At day 10 since hatching, feather tracks started to emerge at the dorsal, wings, and head tracts, while their bills became darker towards the tip (Fig. 3B). By day 13 since hatching, some feathers started unsheathing and presented white tips, while some creamy feathers in their dorsum started to emerge (Fig. 3C). By day 16, larger feather pins were present in their heads and wings, showing a black base and whitish tips; moreover, creamy feathers had fully unsheathed in the dorsal tract (Fig. 3D). By day 22, feathers had fully unsheathed, and the nestlings were fully covered with feathers that were dark in the base and creamy at tip; flight feathers were dark green; the upper mandible was darker; a vaguely defined white postocular spot was present (Fig. 3E). By day 28, the fledglings were observed performing short flights outside the nest; they both conserved the neossotiles in the dorsum, and the general color of the plumage resembled the dark green color of adults, although they did not present any iridescence on the throat; a well-defined postocular white spot was clearly visible at this age (Fig. 3F). In Table 2 we present a summary of the general feather development events according to our observation days.

We were able to determine the hatching day in two nests. The total nesting period, from hatching until fledglings abandoned the nests was 28 and 32 days, respectively.

### Behavioral observations

Incubation and nesting attendance were exclusively performed by the female. During the incubation period, females visited the nest 2–6 times/h; in-nest bouts averaged 8.5 min (range: 1.1–13.7 min), while off-nest bouts averaged 4.9 min (range: 0.5–15.6 min). Females visited their nests to feed nestlings with a frequency of 4–5 visits/h, with an average visit length of 0.3 min (range: 0.1–1.2 min).

### Nest success

We were able to monitor the outcome of seven nests. Two nests (28.5%) were successful and produce two fledglings, two nests (28.5%) were partially successful and produce one fledgling, and three nests (43%) lost their two nestlings. Two of these unsuccessful nests were preyed upon during the incubation period, as indicated by egg remains found in the nests, and one was abandoned during the nestling period, with two fledglings found dead in the nest (Table 1).

Table 2: Feather development table for nestlings of Violet-throated Metaltail *Metallura baroni* found along the Cuenca-Molleturo-Naranjal road, Azuay province, Ecuador, in 2019. Records come from one nest, and correspond to the observations of two nestlings. The day column refers to the day after hatching. N, not visible; V, visible below the skin; P, pins above the skin; U, unsheathing; F, fully unsheathed. The two nestlings presented the same timing in their development.

| Day | Head | Dorsum | Wings | Caudal track |
|-----|------|--------|-------|--------------|
| 1   | N    | N      | N     | N            |
| 10  | V    | P      | P     | P            |
| 13  | P    | U      | U     | U            |
| 16  | U    | U      | U     | U            |
| 22  | F    | F      | F     | F            |

## DISCUSSION

This manuscript presents basic breeding information of the Ecuadorian endemic and threatened *M. baroni* in CNP, knowledge that can be used as a basis to expand on studies that explore the factors that determine breeding success in this species. A breeding period between November and April has been suggested for *M. baroni* (Schuchmann, 1999). Even though we did not find any nests in January and February, our observations fall within the time window suggested by Schuchmann (1999). The bird banding program carried out in montane

forest of CNP (Tinoco *et al.*, 2019) can also provide insights into the seasonality of the breeding season for *M. baroni*. This program samples birds in three climatic seasons of the year: rainy season (sampled in March–April), dry season (sampled in June–July), secondary rainy season (sampled in November) (Tinoco *et al.*, 2019). Of the 105 young *M. baroni* individuals captured in 2006–2018 in this banding program, only three were caught during the rainy season (B. Tinoco, unpubl. data). This result agrees with our observation that the nesting season of *M. baroni* coincides with the main rainy season (March–May). Nonetheless, an even sampling effort for nests across the year, and in different habitat types, is necessary to better determine the seasonality in the breeding period of *M. baroni*.

Regarding the general location of nests, nests architecture and nest materials, we can compare our findings with other description of *M. baroni* nests (Schuchmann, 1999), as well as to other *Metallura* species. Embankments, small cavities, and rocky walls are all common nest locations in *Metallura* hummingbirds (Snow, 1980, Matta-Pereira *et al.*, 2019, Mamani-Cabana, 2020). All the nine nests found were open cup-shaped, coinciding with the only other nest description for *M. baroni* (Schuchmann, 1999). Open cup nests have also been described for Tyrian Metaltail *M. tyrianthina* (Londoño *et al.*, 2014, Matta-Pereira *et al.*, 2019, Greeney & Juiña, 2020, Soto-Patiño, 2021) and Black Metaltail *M. phoebe* (Mamani-Cabana, 2020). We could not find previous information about nest dimension for *M. baroni*, but our measurements were similar to those reported for a *M. tyrianthina* nest in Colombia (Soto-Patiño, 2021), a *M. phoebe* nest in Peru (Mamani-Cabana, 2020), and a Coppery Metaltail *M. theresiae* nest in Peru (González, 2020).

The nesting materials found were similar to those used in nests of other *Metallura* species: mosses, twigs, roots, mammal fur, and spiderweb are all materials found in the external chamber of Neblina Metaltail *M. odomae* (Schuchmann, 1999), *M. phoebe* (Mamani-Cabana, 2020) and *M. tyrianthina* (Matta-Pereira, 2019; Greeney & Juiña, 2020, Soto-Patiño, 2021). Feathers are also reported in the inner chamber of *M. phoebe* (Mamani-Cabana 2020), *M. odomae* (Schuchmann, 1999), *M. theresiae* (González, 2020), and other high elevation hummingbirds, including Andean Hillstar *Oreotrochilus estella* (Carpenter, 1976) and Blue-throated Hillstar *O. cyanolaemus* (Molina *et al.*, 2021). Feathers as nesting material are likely being selected for birds in cold, high elevation environments due to their high insulation properties (Schöl & Hille, 2014). Overall, there is still much to learn about the nests of *Metallura* species, but there are many similarities among nests of all *Metallura* species, supporting the phylogenetic conservatism in nest characters in birds (Fang *et al.*, 2018).

The cold temperatures prevalent at high elevations influence several life-history parameters associated with avian reproduction (Perfito *et al.*, 2014; Boyle *et al.*, 2015), including the duration of the nestling period (Bears *et al.*, 2009; Bründl *et al.*, 2020). For instance, birds breeding at high elevations have longer nestling periods than those breeding in the lowlands (Badyaev, 1997; Altamirano *et al.*, 2015), a pattern probably related to a decrease in the energy available for reproduction in the high mountains (Boyle *et al.*, 2015). In correspondence, the nestling period of 28–32 days observed for *M. baroni* is longer than the nestling periods reported for lowland tropical hummingbirds (18–25 days; Fierro-Calderón & Martin, 2007). However, knowledge about the development time of nestlings is limited for tropical birds, and more studies about the reproduction of birds are needed to further understand how life histories can be shaped by environmental gradients.

We could not find studies on nesting success of tropical Andean hummingbirds to compare our observation of a 43% nest failure in *M. baroni* along a main road. However, nesting along roads may reduce nest survival because of road-related kills (Kuitunen *et al.*, 2003; Dietz *et al.*, 2013). Nesting on the side of the Cuenca-Molleturo-Naranjal road could expose adult females and fledglings alike to collisions with vehicles. Indeed, carcasses of *M. baroni* were reported in a bird road-kill study conducted in CNP (Aguilar *et al.*, 2019). The Cuenca-Molleturo-Naranjal road is likely acting as an ecological trap for birds in CNP (Astudillo *et al.*, 2014; Aguilar *et al.*, 2019), where birds are attracted to the sides of this road despite the high mortality risks that these areas represent (Astudillo *et al.*, 2014). The availability of novel nesting sites for *M. baroni* in the roadside embankments is likely attracting females to build nests along the Cuenca-Molleturo-Naranjal road, where nesting success could be affected. Measuring the effects of the road on *M. baroni* nesting success should be a research priority for the species, with important conservation management implications for CNP.

During this study, Blue-mantled Thornbill *Chalcostigma stanleyi* nested in the same roadside embankments used by *M. baroni*. On several occasions we even observed *M. baroni* females chasing away *C. stanleyi* from nesting areas. A limitation of nesting sites for high Andean birds can promote competitive interactions among

species (Carpenter, 1976), so this possibility should be further evaluated to learn more about interspecific interactions among birds in páramo habitats.

Future studies should expand knowledge about the breeding biology of *M. baroni*, since our results come from data gathered in a single páramo area along a main road. Populations of *M. baroni* are expected to keep declining in the future (BirdLife International, 2021), a problem that could be further aggravated by climate change in the high Andes (Laurance *et al.*, 2011; Freeman *et al.*, 2018). Therefore, sound conservation actions for this species should be guided by scientific knowledge about its natural history and its ecological needs.

#### ACKNOWLEDGEMENTS

We thank Ibeth Alarcón for her support during field activities. BTA was funded by Vicerrectorado de Investigaciones, Universidad del Azuay, grant number 2019-96. This work was conducted under activities related to the permit number 210-2019-DPAA-MA issued by the Ministerio del Ambiente, Agua y Transición Ecológica.



Figure 3: Nestlings' development in a Violet-throated Metaltail *Metallura baroni* nest. A) day 1 (identified by the presence of the remaining egg shale); B) day 10; C) day 13; D) day 16; E) day 22; F) Fledgling outside the nest (Paúl Molina).

## REFERENCES

- Aguilar, J. M., Nieto, A., Espinoza, N., Loja, G., & Tinoco, B. A. (2019). Assessing patterns of bird roadkills in a high Andean Ecuadorian national park. *Studies on Neotropical Fauna and Environment*, 54 (3), 149–156. DOI: <http://doi.org/10.1080/01650521.2019.1649953>
- Altamirano, T. A., Ibarra, J. T., de la Maza, M., Navarrete, S. A., & Bonacic, C. (2015). Reproductive life-history variation in a secondary cavity-nester across an elevational gradient in Andean temperate ecosystems. *The Auk*, 132 (4), 826–835. DOI: <https://doi.org/10.1642/AUK-15-28.1>
- Astudillo, P. X., Samaniego, G. M., Machado, P. J., Aguilar, J. M., Tinoco, B. A., Graham, C. H., Latta, S. & Farwig, N. (2014). The impact of roads on the avifauna of páramo grasslands in Cajas National Park, Ecuador. *Studies on Neotropical Fauna and Environment*, 49 (3), 204–212. DOI: <http://doi.org/10.1080/01650521.2014.960778>
- Astudillo, P. X., Tinoco, B. A., & Siddons, D. (2015). The avifauna of Cajas National Park and Mazán Reserve, southern Ecuador, with notes on new records. *Cotinga*, 37, 1–11. URL: <https://www.neotropicalbirdclub.org/wp-content/uploads/2020/06/C37-Astudillo-et-al.pdf>
- Badyaev, A. V. (1997). Avian life history variation along altitudinal gradients: an example with cardueline finches. *Oecologia*, 111(3), 365–374. DOI: <https://doi.org/10.1007/s004420050247>
- Bears, H., Martin, K., & White, G. C. (2009). Breeding in high-elevation habitat results in shift to slower life-history strategy within a single species. *Journal of Animal Ecology*, 78 (2), 365–375. DOI: <https://doi.org/10.1111/j.1365-2656.2008.01491.x>
- Benham, P. M., Cuervo, A. M., McGuire, J. A., & Witt, C. C. (2015). Biogeography of the Andean metalltail hummingbirds: contrasting evolutionary histories of tree line and habitat-generalist clades. *Journal of Biogeography*, 42 (2), 763–777. DOI: <http://doi.org/10.1111/jbi.12452>
- BirdLife International. (2021). Species factsheet: *Metallura baroni*. BirdLife International. URL: <http://datazone.birdlife.org/species/factsheet/violet-throated-metaltail-metallura-baroni>
- Boyle, A. W., Sandercock, B. K., & Martin, K. (2016). Patterns and drivers of intraspecific variation in avian life history along elevational gradients: a meta-analysis. *Biological Reviews*, 91(2), 469–482. DOI: <https://doi.org/10.1111/brv.12180>
- Bründl, A. C., Sallé, L., Lejeune, L. A., Sorato, E., Thiney, A. C., Chaine, A. S., & Russell, A. F. (2020). Elevational gradients as a model for understanding associations among temperature, breeding phenology and success. *Frontiers in Ecology and Evolution*, 8. URL: <https://www.frontiersin.org/article/10.3389/fevo.2020.563377>
- Campozano, L., Célleri, R., Trachte, K., Bendix, J., & Samaniego, E. (2016) Rainfall and cloud dynamics in the Andes: a southern Ecuador case study. *Advances in Meteorology*, 2016, e3192765. DOI: <http://doi.org/10.1155/2016/3192765>
- Carpenter, F. L. (1976). Ecology and evolution of an Andean hummingbird (*Oreotrochilus estella*). *University of California Publications in Zoology*, 106, 1–74.
- Célleri, R., Willems, P., Buytaert, W., & Feyen, J. (2007). Space-time rainfall variability in the Paute basin, Ecuadorian Andes. *Hydrological Processes*, 21(24), 3316–3327. DOI: <http://doi.org/10.1002/hyp.6575>
- Dietz, M. S., Murdock, C. C., Romero, L. M., Ozgul, A., & Foutopoulos, J. (2013). Distance to a road is associated with reproductive success and physiological stress response in a migratory landbird. *The Wilson Journal of Ornithology*, 125(1), 50–61. DOI: <http://doi.org/10.1676/11-201.1>



- Fang, Y.-T., Tuanmu, M.-N., & Hung C.-M. (2018). Asynchronous evolution of interdependent nest characters across the avian phylogeny. *Nature Communications*, 9, 1863. DOI: <http://doi.org/10.1038/s41467-018-04265-x>
- Fierro-Calderón, K., & Martin, T. E. (2007). Reproductive biology of the Violet-chested Hummingbird in Venezuela and comparisons with other tropical and temperate hummingbirds. *The Condor*, 109(3), 680–685. DOI: <http://doi.org/10.1093/condor/109.3.680>
- Freeman, B. G., Scholer, M. N., Ruiz-Gutiérrez, V., & Fitzpatrick, J. W. (2018). Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences*, 115(47), 11982–11987. DOI: <http://doi.org/10.1073/pnas.1804224115>
- Freile, J. F., & Santander, T. (2005). Áreas importantes para la conservación de las aves en Ecuador. En Boyla, K. A. & A. Estrada (Eds), *Áreas importantes para la conservación de las aves en los Andes Tropicales: sitios prioritarios para la conservación de la biodiversidad* (pp. 283–470). Quito, Ecuador: BirdLife International Conservation Series 14.
- Freile, J. F., Santander G., T., Jiménez-Uzcátegui, G., Carrasco, L., Cisneros-Heredia, D. F., Guevara, E. A., Sánchez-Nivicela, M., & Tinoco, B. A. (2019). *Lista roja de las aves del Ecuador*. Quito, Ecuador: Ministerio del Ambiente, Aves y Conservación, Comité Ecuatoriano de Registros Ornitológicos, Fundación Charles Darwin, Universidad del Azuay, Red Aves Ecuador & Universidad San Francisco de Quito.
- García-Moreno, J., Arcander, P., & Fjeldså, J. (1999). Strong diversification at the treeline among *Metallura* hummingbirds. *The Auk*, 116(3), 702–711. DOI: <https://doi.org/10.2307/4089331>
- González, O. (2020). First description of the nest and juveniles of the Coppery Metaltail (*Metallura theresiae*), with comments on hummingbird nestling obesity. *Wilson Journal of Ornithology*, 132(4), 984–990. DOI: <https://doi.org/10.1676/1559-4491-132.4.984>
- Graves, G. R. (1980). A new species of Metaltail hummingbird from northern Peru. *The Wilson Bulletin*, 92(1), 1–7. URL: <https://sora.unm.edu/sites/default/files/journals/wilson/v092n01/p0001-p0007.pdf>
- Greeney, H. F., & Juíña-J, M. E. (2020). Notas de reproducción de seis especies de colibríes de los Andes en Ecuador. *Ornitología Colombiana*, 18, 1–10. URL: <https://asociacioncolombianadeornitologia.org/ojs/index.php/roc/article/view/414>
- Jongsomjit, D., Jones, S. L., Gardali, T., Geupel, G. R., & Gouse, P. J. (2007). *A guide to nestling development and aging in altricial passerines*. Washington, D.C.: U.S. Department of Interior, Fish and Wildlife Service, Biological Technical Publication, FWS/BTP-R6008- 2007.
- Kuitunen, M. T., Viljanen, J., Rossi, E., & Stenroos, A. (2003). Impact of busy roads on breeding success in Pied Flycatchers *Ficedula hypoleuca*. *Environmental Management*, 31(1), 79–85. DOI: <https://doi.org/10.1007/s00267-002-2694-7>
- Laurance, W. F., Carolina Useche, D., Shoo, L. P., Herzog, S. K., Kessler, M., Escobar, F., Brehm, G., Axmacher, J. C., Chen, I.-C., Gámez, L. A., Hietz, P., Fiedler, K., Pyrcz, T., Wolf, J., Merkord, C. L., Cardelus, C., Marshall, A. R., Ah-Peng, C., Aplet, G. H., Arizmendi, M. C., Baker, W. J., Barone, J., Brühl, C. A., Bussmann, R. W., Cicuzza, D., Eilu, G., Favila, M. E., Hemp, A., Hemp, C., Homeier, J., Hurtado, J., Jankowski, J., Kattan, G., Kluge, J., Krömer, T., Lees, D. C., Lehnert, M., Longino, J. T., Lovett, J., Martin, P. H., Patterson, B. C., Pearson, R. G., Peh, K. S. H., Richardson, B., Richardson, M., Samways, M. J., Senbeta, F., Smith, T. B., Utteridge, T. M. A., Watkins, J. E., Wilson, R., Williams, S. E., & Thomas, C. D. (2011). Global warming, elevational ranges and the vulnerability of tropical biota. *Biological Conservation*, 144(1), 548–557. DOI: <https://doi.org/10.1016/j.biocon.2010.10.010>
- Londoño, G. A. (2014). Anidación de aves en un gradiente altitudinal. Parque Nacional del Manu, Perú. Rapid Color Guide #514. Chicago, IL: The Field Museum. URL: <http://fieldguides.fieldmuseum.org/guides/guide/514>

- Mamani-Cabana, W, N. (2020). Description of the nest and eggs of the Black Metaltail (*Metallura phoebe*) in high Andean *Polylepis* forest. *The Wilson Journal of Ornithology*, 131(4), 825–828. DOI: <https://doi.org/10.1676/1559-4491-131.4.825>
- Matta-Pereira, M, E., Lentino, M., & Muñoz, D. (2019). El nido de Metalura Tiria *Metallura tyrianthina* en Venezuela y Perú. *Cotinga*, 41, 24–28.
- Molina, P., Pacheco, D., Carrasco A., & Tinoco, B. A. (2021). On the breeding biology of the Blue-throated Hillstar (*Oreotrochilus cyanolaemus*), a recently discovered species from Southern Ecuador. *The Wilson Journal of Ornithology*, 132(4), 1007–1014. DOI: <https://doi.org/10.1676/1559-4491-132.4.1007>
- Perfito, N., Tramontin, A. D., Meddle, S., Sharp, P., Afik, D., Gee, J., Ishii, S., Kikuchi, M., & Wingfield, J. C. (2004). Reproductive development according to elevation in a seasonally breeding male songbird. *Oecologia*, 140(2), 201–210. DOI: <https://doi.org/10.1007/s00442-004-1576-5>
- Rodríguez-Girón, S., Rodas, F., Schubert, A., & Vasco Tapia, S. (2014). *Área de Biósfera Macizo del Cajas, experiencias de desarrollo sostenible para el bien vivir*. Cuenca, Ecuador: ETAPA EP, Municipio de Cuenca, Cooperación Alemana GIZ & Naturaleza y Cultura Internacional.
- Schöl, E. M., & Hille, S. M. (2014). Do Great Tits *Parus major* nesting at high altitudes build better insulated eggs? *Ardeola*, 61(2), 323–333. DOI: <https://doi.org/10.13157/arla.61.2.2014.323>
- Schuchmann, K. L. (1999). Family Trochilidae (Hummingbirds). In del Hoyo, J., Elliott, A., & J. Sargatal (Eds), *Handbook of the birds of the world. Volume 5. Barn-owls to hummingbirds* (pp 468–680). Barcelona, Spain: Lynx Edicions.
- Simon, J., & Pacheco, S. (2005). On the standardization of nest descriptions of neotropical birds. *Revista Brasileira de Ornitologia*, 13(23), 143–154. URL: <http://revbrasilornitol.com.br/BJO/article/view/2201>
- Soto-Patiño, J., Pérez-Peña, S., Córdoba-Córdoba, S., & Ocampo, D. (2021). On the nests, eggs, and reproductive evidence of Tyrian Metaltail (*Metallura tyrianthina*) and Glowing Puffleg (*Eriocnemis vestita*) in the northeastern Colombian Andes. *The Wilson Journal of Ornithology*, 133(1), 110–116. DOI: <https://doi.org/10.1676/19-00147>
- Tinoco, B. A., Astudillo, P. X., Latta, S. & Graham, C. H. (2009). Distribution, ecology and conservation of an endangered Andean hummingbird: The Violet-throated Metaltail (*Metallura baroni*). *Bird Conservation International*, 19(1), 63–76. DOI: <https://doi.org/10.1017/S0959270908007703>
- Tinoco, B. A, Graham, L., Astudillo, P. X., Nieto, A., Aguilar, J. M., Latta, S. C. & Graham, C. H. (2019). Survival estimates of bird species across altered habitats in the tropical Andes. *Journal of Field Ornithology*, 90(2), 105–116. DOI: <https://doi.org/10.1111/jofo.12293>
- Zyskowski, K., & Prum, R. O. (1999). Phylogenetic analysis of the nest architecture of Neotropical ovenbirds (Furnariidae). *The Auk*, 116(4), 891–911. DOI: <https://doi.org/10.2307/4089670>